





DRUG TEST

Self-test for the detection of 6 drugs in urine: Morphine, Amphetamine, Cocaine, Marijuana, Methamphetamine and Methadone



DETECTED SUBSTANCES

TEST	Cut-off (ng/mL)	Time permanence in urine after exposure	
Amphetamine (AMP1000)	1000	1-3 days for a single administration, up to 9 days in chronic users	
Cocaine (COC 300)	300	1-3 days for a single administration, up to 3 weeks in chronic users	
Marijuana	50	single administration (smoking): 30 h on average, up to 4 days single administration (oral intake): up to 6 days occasional use (once or twice a week): up to 30 days regular consumption: up to 3 months	
Methadone (MTD 300)	300	1,5 to 3 days	
Methamphetamine (MET 500)	500	1 to 4 days for a single administration, up to 5 days in chronic users	
Morphine (MOP 300)	300	10-55 h for a single administration, up to 11 days in chronic users	

WHO ARE THE INTENDED USERS

DRUG TEST can be used by everyone who wants to check the presence of the above mentioned substances in urine.

TEST PRINCIPLE

DRUG TEST is an immunochromatographic assay that can be performed without the use of an instrument. The test utilizes monoclonal antibodies to selectively detect levels of specific drugs in urine.

TECH SPECS

	CUT-OFF	SENSITIVITY	SPECIFICITY	OVERALL ACCURACY
AMP	1000 ng/mL	99,90%	99,90%	99,90%
MET	500 ng/mL	99,90%	99,90%	99,90%
COC	300 ng/mL	99,90%	99,90%	99,90%
MOP	300 ng/mL	99,90%	99,90%	99,90%
MTD	300 ng/mL	99,90%	99,90%	99,90%
ТНС	50 ng/mL	99,09%	99,90%	99,90%

HOW TO USE IT

1) Open the cap, with the arrow pointing toward the urine specimen, dip the test panel vertically in the urine specimen for at least 10 to 15 seconds. 2) Replace the cap and leave the test panel on a non-absorbent flat surface. 3) Wait 5 minutes before reading the result.



CONTENT: 1 sealed aluminium pouch containing: 1 test panel and 1 desiccant bag; 1 instructions for use leaflet.

\triangleright	VIDEO TUTORIAL	REF DESCRIPTION		NUMBER OF TESTS	SHELF LIFE
		19806-1	DRUG TEST	1 TEST	24 MONTHS

